arXiv:0806.2270v2 [hep-th] 20 Jun 2008

An $\mathcal{N} = 1$ Superfield Action for M2 branes

Andrea Mauri[1*] **and Anastasios C. Petkou**[1†]

[1] *Department of Physics, University of Crete, Heraklion 71003, Greece*

Abstract

We present an octonionic $\mathcal{N} = 1$ superfield action that reproduces in components the action of Bagger and Lambert for M2 branes. By giving an expectation value to one of the scalars we obtain the maximally supersymmetric superfield action for D2 branes.

1 Introduction

Despite its importance for M-theory and higher-spin gauge theory holography[1] AdS$_4$/CFT$_3$ correspondence is essentially unexplored in comparison to its AdS$_5$/CFT$_4$ counterpart. Presumably, the main reason has been the lack so far of a manifestly maximally superconformal invariant 2+1 dimensional theory similar in status to $\mathcal{N} = 4$ SYM in 3+1 dimensions. It seems possible that this obstacle has been overcome by the proposal of Bagger and Lambert [2, 3, 4, 5] of a $\mathcal{N} = 8$ superconformal theory in 2+1 dimensions, with a non-standard gauge structure based on 3-algebras. This theory has attracted very much interest in the past few months and many interesting results have appeared. Multiple M2 branes have been discussed in [7, 9, 10, 11, 13, 14, 16, 28, 8, 41, 17, 30]. The relation of M2 to D2 branes was elucidated in [7, 12, 23, 25, 26, 43]. Algebraic aspects of 3-algebras have been discussed in [6, 15, 19, 22, 24, 33, 37, 38, 40]. General aspects of three-dimensional Chern-Simons theories and extensions have been discussed in [36, 20] and also in [29, 31, 18, 21, 27, 32, 35, 34]. Very recently, an interesting class of $U(N) \times U(N)$ Chern-Simons theories that may describe multiple M2 branes have been discussed in [39, 41, 42].

In this short note, we present an $\mathcal{N} = 1$ superfield action whose component expansion gives the BL theory for a 3-algebra with totally antisymmetric structure constants f^{abcd}. We use real three-dimensional superfields both for the matter as well as for the Chern-Simons part of the action. The crucial point is the use of the octonionic self-dual tensor in the construction of the real superpotential. In this way, the superpotential is only manifestly $SO(7)$ invariant. However, for specially chosen couplings, the component action coincides with the BL action, and hence full $SO(8)$ symmetry is restored. We believe that octonions will play a fundamental role in future studies of AdS$_4$/CFT$_3$.

---

[*] Andrea.Mauri@mi.infn.it

[†] petkou@physics.uoc.gr

[1] For a general discussion see [1].

Our motivation comes in part from corresponding studies in AdS$_5$/CFT$_4$ where the $\mathcal{N} = 1$ formulation of $\mathcal{N} = 4$ SYM has been an extremely efficient tool for studies of anomalous dimensions, non-renormalization properties and integrability. We believe that our $\mathcal{N} = 1$ action will be similarly useful in this case too.

As a simple test for our action we follow [7] and demonstrate that giving an expectation value in one of the scalar supefields, our action yields the maximally supersymmetric YM theory in 2+1 dimensions, as it should.

2 The $\mathcal{N} = 1$ superfield action

We consider eight real $\mathcal{N} = 1$ superfields[2] as

$$\Phi^I_a = \phi^I_a + \theta^{\alpha,\dot{8}}\hat{\Gamma}^I_{\dot{8}A}\psi^A_{\alpha,a} - \theta^2 F^I_a\,,\quad I, A = 1, 2, .., 8. \tag{1}$$

where a denotes the index of the three-algebra algebra with structure constants f^{abcd}. We use the $SO(8)$ triality tensor $\Gamma^I_{A\dot{A}}$, $I, \dot{A}, A = 1, 2, .., 8$. In the representation where $\hat{\Gamma}^I_{\dot{8}A} = -\delta^I_A$ (see appendix B), choosing the superspace coordinate to point in the $\dot{8}$ direction we have essentially made equivalent the vector and one of the two spinorial representations of $SO(8)$. We use the notation of [46] summarized along with other useful relations and conventions in the Appendix A.

Our superfield action couples the matter superfields to a Chern-Simons gauge superfield Γ^α_{ab} in the Wess-Zumino gauge, that carries two gauge group indices a, b

$$\begin{aligned} S \;=\; \int d^3x\, d^2\theta \Big[& 2\gamma \left(D^\alpha \Phi^I_d - f^{abc}{}_d\, \Gamma^\alpha{}_{ab}\Phi^I_c\right)^2 + \alpha\, f^{abcd}\,(D^\alpha\Gamma^\beta{}_{ab})(D_\beta\Gamma_{\alpha\, cd}) + \\ & +\beta\, f^{cda}{}_g f^{efgb}\,(D^\alpha\Gamma^\beta{}_{ab})\Gamma_{\alpha\, cd}\Gamma_{\beta\, ef} + k\, f^{abcd}\, C_{IJKL}\,\Phi^I_a\Phi^J_b\Phi^K_c\Phi^L_d \Big]\,. \end{aligned} \tag{2}$$

Apart from the overall normalization, the only adjustable parameter is the coupling constant of the real superpotential. Nevertheless, we keep all coefficients α, β, γ, k arbitrary having in mind possible generalizations of the action (2).

The crucial point is the use of the self-dual[3] eight dimensional tensor C_{IJKL}, $I, J, K, L = 1, 2, ..8$ that describes the embeddings of $SO(7)$ into $SO(8)$ [47, 48, 49, 50]. Its properties are briefly recalled in the Appendix B. Hence, the presence of the superpotential implies that (2) has only $SO(7)$ manifest global symmetry. Our strategy is to fix the coefficients in (2) by comparing the resulting component action with the one of BL. In this way, the $SO(7)$ symmetry is enhanced to $SO(8)$ and hence we achieve maximal supersymmetry.

We detail next the various projections.

A) *Scalar kinetic terms*

$$D^2 \left[2\gamma \left(D^\alpha \Phi^I_d - f^{abc}{}_d\, \Gamma^\alpha{}_{ab}\Phi^I_c\right)^2\right]\Big|_{\theta=0} =$$

---

[2] Earlier works on the superfield formulation of Chern-Simons theories coupled to matter include [44, 45].

[3] Similar result can be obtained using the anti self-dual tensor.

$$
\begin{aligned}
&= -2\gamma\, F^I_{\ d} F^{I\,d} - 2\gamma\, \phi^I_{\ d}\, \Box\, \phi^{I\,d} - 2\gamma\, i\, \psi^{\alpha\, I}_{\ d} \partial_\alpha^{\ \beta} \psi_\beta^{\ I\,d} + 2\gamma f^{abc}_{\ \ \ d} f^{efgd}\, A^\mu_{\ ab} A_{\mu\, ef}\, \phi^I_{\ c} \phi^I_{\ g} - \\
&\quad -\gamma\, f^{abc}_{\ \ \ d} \Big[-\phi^I_{\ c}\, (\gamma_\mu)^\alpha_{\ \gamma}\, A^\mu_{\ ab} \partial^\gamma_{\ \alpha}\, \phi^{I\,d} - \left(\partial^\alpha_{\ \gamma} \phi^{I\,d}\right) (\gamma_\mu)^\gamma_{\ \alpha}\, A^\mu_{\ ab}\, \phi^I_{\ c} + \\
&\qquad\qquad +2i\, \psi^{\gamma\, I}_{\ c}\, (\gamma_\mu)_\gamma^{\ \alpha}\, A^\mu_{\ ab} \psi^{I\,d}_\alpha + 4\lambda^\alpha_{\ ab}\, \psi^{I\,d}_\alpha \phi^I_{\ c} \Big] = \\
&= 2\gamma \Big(\nabla^\mu \phi^I_{\ d}\Big) \Big(\nabla_\mu \phi^{I\,d}\Big) - 2i\, \gamma\, \psi^{\alpha\, I}_{\ d} \nabla_\alpha^{\ \beta}\, \psi^{I\,d}_\beta - 2\gamma\, F^I_{\ d} F^{I\,d} - 4\gamma f^{abc}_{\ \ \ d} \lambda^\alpha_{\ ab}\, \psi^{I\,d}_\alpha \phi^I_{\ c}\,,
\end{aligned}
\tag{3}
$$

where

$$
\nabla_\mu \phi^I_d \equiv \partial_\mu \phi^{I\,d} - \ f^{abc}_{\ \ \ d}\, A_{\mu\, ab}\, \phi^{I\,c}\,. \tag{4}
$$

B) *Chern-Simons terms*:

$$
\begin{aligned}
&D^2 \left[\alpha\, f^{abcd}\, (D^\alpha \Gamma^\beta_{\ ab})(D_\beta \Gamma_{\alpha\, cd}) + \ \beta\, f^{cda}_{\ \ \ g} f^{efgb}\, (D^\alpha \Gamma^\beta_{\ ab}) \Gamma_{\alpha\, cd} \Gamma_{\beta\, ef} \right] \Big|_{\theta=0} = \\
&= 4\, \alpha\, f^{abcd} \lambda^\alpha_{\ ab} \lambda_{\alpha\, cd} - 4\alpha\, f^{abcd}\, \epsilon^{\mu\nu\rho}\, A_{\mu\, ab}\, \partial_\nu A_{\rho\, cd} - 2\beta\, f^{cda}_{\ \ \ g} f^{efgb}\, \epsilon^{\mu\nu\rho}\, A_{\mu\, ab}\, A_{\nu\, cd}\, A_{\rho\, ef}\,.
\end{aligned}
\tag{5}
$$

C) *Superpotential*:

$$
D^2 \Big[k\, f^{abcd}\, C_{IJKL}\, \Phi^I_{\ a} \Phi^J_{\ b} \Phi^K_{\ c} \Phi^L_{\ d} \Big] \Big|_{\theta=0} = k\, f^{abcd}\, C_{IJKL} \Big[6\, \psi^{\alpha I}_{\ a} \psi^J_{\alpha\, b} \phi^K_{\ c} \phi^L_{\ d} + 4\, F^I_{\ a} \phi^J_{\ b} \phi^K_{\ c} \phi^L_{\ d} \Big]\,. \tag{6}
$$

The equations of motion for the auxiliary fields read

$$
\begin{aligned}
F^{Ia} &= \frac{k}{\gamma}\, f^{abcd}\, C^I_{\ JKL}\, \phi^J_{\ b} \phi^K_{\ c} \phi^L_{\ d}\,, &(7)\\
\lambda_{\alpha\ ab} &= \frac{\gamma}{2\alpha} \phi^I_{\ a} \psi^I_{\alpha\, b}\,. &(8)
\end{aligned}
$$

Using (7) and (8), the terms contributing to the potential from A), B) and C) that involve the auxiliary fields give

$$
\begin{aligned}
&-\frac{\gamma^2}{\alpha} f^{abcd} \psi^{\alpha\, I}_a \phi^I_{\ b} \psi^J_{\alpha\, c} \phi^J_{\ d} + \frac{2k^2}{\gamma}\, f^{bcda}\, f^{efg}_{\ \ \ \ a}\, C^I_{\ OMN}\, C_{IJKL}\, \phi^O_{\ e} \phi^M_{\ f} \phi^N_{\ g} \phi^J_{\ b} \phi^K_{\ c} \phi^L_{\ d} = \\
&\qquad -\frac{\gamma^2}{\alpha} f^{abcd} \psi^{\alpha\, I}_a \phi^I_{\ b} \psi^J_{\alpha\, c} \phi^J_{\ d} + \frac{12k^2}{\gamma}\, f^{bcda}\, f^{efg}_{\ \ \ \ a}\, \phi^J_{\ e} \phi^K_{\ f} \phi^L_{\ g} \phi^J_{\ b} \phi^K_{\ c} \phi^L_{\ d} + \\
&\qquad\qquad - \frac{18k^2}{\gamma}\, f^{bcda}\, f^{efg}_{\ \ \ \ a}\, C_{KLOM}\, \phi^J_{\ e} \phi^K_{\ f}\, \phi^L_{\ g} \phi^O_{\ b} \phi^M_{\ c}\, \phi^J_{\ d} = \\
&= \qquad -\frac{\gamma^2}{\alpha} f^{abcd} \psi^{\alpha\, I}_a \phi^I_{\ b} \psi^J_{\alpha\, c} \phi^J_{\ d} + \frac{12k^2}{\gamma}\, \mathrm{Tr}\Big([\phi^J, \phi^K, \phi^L], [\phi^J, \phi^K, \phi^L] \Big)\,.
\end{aligned}
\tag{9}
$$

To arrive at the last line we have used the contraction of the self-dual form [47, 48, 50]

$$
C^I_{\ OMN}\, C^{IJKL} = \delta^{[J}_O \delta^{\ K}_M \delta^{L]}_N - 9\, C^{KL}_{\ \ \ OM} \delta^J_N\,. \tag{10}
$$

It is also crucial that the third line in (9) vanishes. This can be shown for a general 3-algebra with totally antisymmetric structure constants f^{abcd}. Indeed, using the fundamental identity

$$
f^{bcda}\, f^{efg}_{\ \ \ \ a} = f^{efda}\, f^{bcg}_{\ \ \ \ a} + f^{efba}\, f^{cdg}_{\ \ \ \ a} + f^{efca}\, f^{dbg}_{\ \ \ \ a} \tag{11}
$$

in the relevant part of the third line of (9) we obtain after some relabeling

$$\begin{aligned} f^{bcda}\, f^{efg}{}_{a}\, C_{KLOM}\, \phi^{J}{}_{e} \phi^{K}_{f} \phi^{L}_{g} \phi^{O}_{b} \phi^{M}_{c} \phi^{J}_{d} &= -2 f^{bcga}\, f^{efd}{}_{a} C_{KLOM} \phi^{J}_{e} \phi^{K}_{f} \phi^{L}_{g} \phi^{O}_{b} \phi^{M}_{d} \phi^{J}_{c} \\ &= -2 f^{bcda}\, f^{efg}{}_{a} C_{KLOM} \phi^{J}_{e} \phi^{K}_{f} \phi^{L}_{g} \phi^{O}_{b} \phi^{M}_{c} \phi^{J}_{d}\,, \end{aligned} \tag{12}$$

where we have also used the antisymmetry of C_{KLOM}. Hence this term vanishes. Putting everything together we arrive at the component action

$$\begin{aligned} S \;=\; \int d^3x \Big[& 2\gamma \Big(\nabla^{\mu} \phi^{I}{}_{d}\Big) \Big(\nabla_{\mu} \phi^{I\,d}\Big) - 2\gamma i \psi^{\alpha\,I}_{\;d} \nabla_{\alpha}^{\;\beta}\, \psi^{I\,d}_{\beta} + \\ & -4\,\alpha\, f^{abcd}\, \epsilon^{\mu\nu\rho}\, A_{\mu\,ab}\, \partial_{\nu} A_{\rho\,cd} - 2\beta\, f^{cda}{}_{g} f^{efgb}\, \epsilon^{\mu\nu\rho}\, A_{\mu\,ab}\, A_{\nu\,cd}\, A_{\rho\,ef} + \\ & +6\,k\, f^{abcd} \,(C_{IJKL} + \frac{\gamma^2}{6k\alpha} \delta_{IK} \delta_{JL})\, \psi^{\alpha I}_{\;a} \psi^{J}_{\alpha\,b} \phi^{K}_{\;c} \phi^{L}_{\;d} + \frac{12k^2}{\gamma} \mathrm{Tr}\Big([\phi^I, \phi^J, \phi^K], [\phi^I, \phi^J, \phi^K] \Big) \Big]\,. \end{aligned} \tag{13}$$

Now, in our representation we have

$$\psi^{\alpha\,I}_{a} = \hat{\Gamma}^{I}_{\dot{8}A} \psi^{\alpha\,A}_{a} = -\delta^{I}_{A} \psi^{\alpha\,A}_{a}\,, \tag{14}$$

so that for the spinor kinetic term we have

$$-\,2\gamma i \psi^{\alpha\,I}_{\;d} \nabla_{\alpha}^{\;\beta}\, \psi^{I\,d}_{\beta} = -2\gamma i \psi^{\alpha\,A}_{\;d} \nabla_{\alpha}^{\;\beta}\, \psi^{A\,d}_{\beta}\,. \tag{15}$$

Then, choosing

$$\alpha = -\frac{1}{8} \qquad \beta = -\frac{1}{6} \qquad \gamma = -\frac{1}{4} \qquad k = -\frac{1}{24}\,, \tag{16}$$

we obtain the action

$$\begin{aligned} S \;=\; \int d^3x \Big[& -\frac{1}{2} \Big(\nabla^{\mu} \phi^{I}{}_{d}\Big) \Big(\nabla_{\mu} \phi^{I\,d}\Big) + \frac{i}{2} \psi^{\alpha\,A}_{\;d} \nabla_{\alpha}^{\;\beta}\, \psi^{A\,d}_{\beta} + \\ & +\frac{1}{2} f^{abcd}\, \epsilon^{\mu\nu\rho}\, A_{\mu\,ab}\, \partial_{\nu} A_{\rho\,cd} + \frac{1}{3} f^{cda}{}_{g} f^{efgb}\, \epsilon^{\mu\nu\rho}\, A_{\mu\,ab}\, A_{\nu\,cd}\, A_{\rho\,ef} + \\ & -\frac{1}{4}\, f^{abcd} \,(C_{IJKL} + \delta_{IK} \delta_{JL} - \delta_{IL} \delta_{JK})\, \psi^{\alpha I}_{\;a} \psi^{J}_{\alpha\,b} \phi^{K}_{\;c} \phi^{L}_{\;d} - \frac{1}{12} \mathrm{Tr}\Big([\phi^I, \phi^J, \phi^K], [\phi^I, \phi^J, \phi^K] \Big)\,. \end{aligned} \tag{17}$$

This coincides exactly with the Bagger and Lambert action given in [3]. To see that, notice that in our notations we use the purely imaginary charge conjugation matrix C to raise and lower spinor indices. Therefore for a real Majorana spinor ψ we have the identifications

$$(\bar{\psi})^{\alpha} = (\psi^{T} C)^{\alpha} = C^{\alpha\beta} \psi_{\beta} = \psi^{\alpha} = i (\bar{\psi}_{BL})^{\alpha}\,, \tag{18}$$

and also to match the fermion kinetic term

$$i\, \psi^{\alpha} (\gamma^{\mu} \partial_{\mu})_{\alpha}^{\;\beta}\, \psi_{\beta} = \left(\bar{\psi}_{BL}\right)^{\alpha} (\gamma^{\mu} \partial_{\mu})_{\alpha}^{\;\beta}\, (\psi_{BL})_{\beta} = i\, \left(\bar{\psi}_{BL}\right)^{\alpha} (\gamma^{\mu}_{BL} \partial_{\mu})_{\alpha}^{\;\beta}\, (\psi_{BL})_{\beta}\,, \tag{19}$$

that requires (note the position of the γ-matrix indices)

$$(\gamma^{\mu}_{BL})_{\alpha}^{\;\beta} = i\, (\gamma^{\mu})_{\alpha}^{\;\beta}\,. \tag{20}$$

Our final superspace action is then written as

$$\begin{aligned} S &= \int d^3x\, d^2\theta \Big[-\frac{1}{2}\left(D^\alpha \Phi^I_d - f^{abc}{}_d\, \Gamma^\alpha{}_{ab} \Phi^I_c\right)^2 - \frac{1}{8} f^{abcd}\,(D^\alpha \Gamma^\beta{}_{ab})(D_\beta \Gamma_{\alpha\, cd}) + \\ & -\frac{1}{6} f^{cda}{}_g f^{efgb}\,(D^\alpha \Gamma^\beta{}_{ab})\Gamma_{\alpha\, cd}\Gamma_{\beta\, ef} - \frac{1}{24} f^{abcd}\, C_{IJKL}\, \Phi^I_a \Phi^J_b \Phi^K_c \Phi^L_d \Big]\,. \end{aligned} \tag{21}$$

3 M2 to D2

As a simple test for our action (21) we give an expectation value to one of the scalars. Following [7] we expect that the resulting action will be the maximally supersymmetric YM theory in 2+1 dimensions for the gauge group $SU(2)$. We split the $I = 1, .., 8$ index of the scalar superfield as $I \mapsto (i, 8)$ with $i = 1, ..., 7$ and the $SO(4)$ index $\hat{a} = 1, ..., 4$ as $\hat{a} \mapsto (a, x)$ with $a = 1, ..., 3$. Then we give an expectation value to the scalar superfield which we identify with the dimensionful coupling constant of the 2+1 SYM as

$$\langle \Phi^8_x \rangle = g_{YM}\,. \tag{22}$$

For the spinor superfield we define

$$\Gamma^\alpha_{ax} \equiv A^\alpha_a\,, \tag{23}$$

$$\epsilon^{abc}\Gamma^\alpha_{ab} \equiv B^\alpha_c\,. \tag{24}$$

We rewrite the superspace action in terms of the new fields and indices.

i) *Kinetic terms*:

$$\begin{aligned} 2\gamma(D^\alpha \Phi^I_{\hat{d}} - \epsilon^{\hat{a}\hat{b}\hat{c}}{}_{\hat{d}}\Gamma^\alpha_{\hat{a}\hat{b}}\Phi^I_{\hat{c}})^2 &= \gamma\left[\nabla^\alpha \Phi^i_d \nabla_\alpha \Phi^{id} + \nabla^\alpha \Phi^8_d \nabla_\alpha \Phi^{8d} + D^\alpha \Phi^i_x D_\alpha \Phi^{ix} + \right. \\ &\left. + B^\alpha_d(2\Phi^i_x \nabla_\alpha \Phi^{id} + 2g_{YM}\nabla_\alpha \Phi^{8d} - 2\Phi^{id}D_\alpha \Phi^{ix}) + B^\alpha_d B^d_\alpha(g^2_{YM} + \Phi^i_x\Phi^i_x) + B^{\alpha c}B^g_\alpha \Phi^i_c \Phi^i_g\right] \end{aligned} \tag{25}$$

where we defined the gauge covariant derivative

$$\nabla^\alpha \Phi^i_d = D^\alpha \Phi^i_d - 2\epsilon^{bcd}A^\alpha_b \phi^i_c \tag{26}$$

ii) *Superpotential term*:

$$k\,\epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}}\,C_{IJKL}\,\Phi^I_{\hat{a}}\Phi^J_{\hat{b}}\Phi^K_{\hat{c}}\Phi^L_{\hat{d}} = 4\,k\,\epsilon^{bcd}\,c_{jkl}\,g_{YM}\,\Phi^j_b\Phi^k_c\Phi^l_d + \cdots\,, \tag{27}$$

where the dots indicate subleading terms in the large g_{YM} limit which we have discarded.

iii) *Chern-Simons terms*:

$$\begin{aligned} &\alpha\,\epsilon^{\hat{a}\hat{b}\hat{c}\hat{d}}\,(D^\alpha\Gamma^\beta{}_{\hat{a}\hat{b}})D_\beta\Gamma_{\alpha\,\hat{c}\hat{d}} + \beta\,\epsilon^{\hat{c}\hat{d}\hat{a}}{}_{\hat{g}}\,\epsilon^{\hat{e}\hat{f}\hat{g}\hat{b}}\,(D^\alpha\Gamma^\beta{}_{\hat{a}\hat{b}})\Gamma_{\alpha\,\hat{c}\hat{d}}\Gamma_{\beta\,\hat{e}\hat{f}} = 4\alpha\,B^{\beta d}(D^\alpha D_\beta A_{\alpha d}) + \\ &+2\beta\,\epsilon^{dag}\,B^\beta_g\left[2(D^\alpha A_{\beta a})A_{\alpha\, d} + (D_\beta A^\alpha_a)A_{\alpha d} + (D^\alpha A_{\alpha a})A_{\beta d}\right] + \frac{\beta}{2}\,\epsilon^{abc}(D^\alpha B^\beta_c)B_{\alpha a}B_{\beta b} = \\ &= B^\beta_d\left[4\alpha\,D^\alpha D_\beta A^d_\alpha + 6\beta\epsilon^{dba}(D^\alpha A_{\beta a})A_{\alpha\, b}\right] + \frac{\beta}{2}\,\epsilon^{abc}(D^\alpha B^\beta_c)B_{\alpha a}B_{\beta b}\,. \end{aligned} \tag{28}$$

To arrive in the the last line we have used the fact that in the Wess-Zumino gauge $D^\alpha \Gamma_\alpha$ is vanishing and $D_\alpha \Gamma_\beta$ can be symmetrized.

Next, we derive the equations of motions for the B auxiliary superfield neglecting the terms cubic in B and also terms of the form $B^2\Phi^2$, as in [7]. We thus obtain

$$B_{\alpha\, d} = -\frac{1}{2\gamma g_{YM}^2}\Big[W_{\alpha\, d} + 2\gamma(\Phi^i_x \nabla_\alpha \Phi^i_d + g_{YM}\nabla_\alpha \Phi^8_d - \Phi^i_d D_\alpha \Phi^{ix})\Big]\,, \tag{29}$$

where we defined

$$W^d_\beta = 4\alpha\, D^\alpha D_\beta A^d_\alpha + 6\beta \epsilon^{dba}(D^\alpha A_{\beta a})A_{\alpha\, b}\,. \tag{30}$$

Inserting B into the action and collecting all the terms we get

$$\begin{aligned} \mathcal{L} \;=\;& -\frac{1}{2\gamma g_{YM}^2}[W^\alpha_d + 2\gamma(\Phi^i_x \nabla^\alpha \Phi^i_d + g_{YM}\nabla^\alpha \Phi^8_d - \Phi^i_d D^\alpha \Phi^{ix})]^2 + 4\,k\,\epsilon^{bcd}\,c_{jkl}\,g_{YM}\,\Phi^j_b\Phi^k_c\Phi^l_d + \\ & +\gamma\left[\nabla^\alpha \Phi^i_d \nabla_\alpha \Phi^{id} + \nabla^\alpha \Phi^8_d \nabla_\alpha \Phi^{8d} + D^\alpha \Phi^i_x D_\alpha \Phi^{ix}\right] = \\ & = \frac{1}{g_{YM}^2} W^\alpha_d W^d_\alpha - \frac{1}{4}\nabla^\alpha \Phi^i_d \nabla_\alpha \Phi^{id} - \frac{1}{4} D^\alpha \Phi^i_x D_\alpha \Phi^{ix} - \frac{1}{6}\,\epsilon^{bcd}\,c_{jkl}\,g_{YM}\,\Phi^j_b\Phi^k_c\Phi^l_d + \ldots\,, \end{aligned} \tag{31}$$

where the dots stand for subleading contributions and we substituded the values in (16), so that now

$$W^d_\beta = \frac{1}{2}\, D^\alpha D_\beta A^d_\alpha + \epsilon^{dba}(D^\alpha A_{\beta a})A_{\alpha\, b}\,. \tag{32}$$

Rescaling the gauge superfield as in [7] $A \rightarrow 1/2A$ we see that $W^\alpha_d W^d_\alpha$ gives the right SYM kinetic term in the Wess-Zumino gauge and the covariant derivative in (26) assumes its standard form. Finally, ignoring the contribution of Φ^i_x that has completely decoupled, we arrive at

$$\mathcal{L} = \frac{1}{8g_{YM}^2}(D^\alpha D_\beta A^d_\alpha + \epsilon^{dba}(D^\alpha A_{\beta a})A_{\alpha\, b})^2 - \frac{1}{4}\nabla^\alpha \Phi^i_d \nabla_\alpha \Phi^{id} - \frac{1}{6}\,\epsilon^{bcd}\,c_{jkl}\,g_{YM}\,\Phi^j_b\Phi^k_c\Phi^l_d\,. \tag{33}$$

This is the superfield Langangian for maximally supersymmetric YM in 2+1 dimensions for $SU(2)$. It is intriguing to notice that the Lagrangian (33) is remarkably similar to the octonionic $\mathcal{N} = (1,1)$ sigma model (in two dimensions) of [51].

4 Conclusions

We have presented an $\mathcal{N} = 1$ superfield action in three dimensions that in components gives the Bagger-Lambert action for a general 3-algebra with totally antisymmetric structure constants f^{abcd}. Crucial in our construction were the self-dual octonionic tensors C_{IJKL}. Although the tensors are $SO(7)$ invariant, we have shown that a special choice of the parameters in the action enhances the global symmetry to $SO(8)$. We have demonstrated that a superhiggs mechanism yields the maximally supersymemtric 2+1 YM theory on D2 branes, curiously in a formalism resembling a two-dimensional sigma model. We hope that our superfield action and its generalizations can be used in $\mathcal{N} = 1$ superfield calculations that should shed more light into the AdS$_4$/CFT$_3$ correspondence.

Acknowledgments

We would like to thank A. Santambrogio for useful discussions. We are also grateful to Hai Lin for very useful correspondence after the first version of the paper. The work of A.C.P. is partially supported by the European RTN Program MRTN-CT-2004-512194.

A Superspace Notations

In this appendix we collect the useful identities for our superfields and gamma matrices. We follow Superspace [46].

The component field definitions are as

$$\begin{array}{rclrcl}
\Phi^I_a| & \equiv & \phi^I_a & D_\alpha\Phi^I_a| & \equiv & \psi^I_{\alpha\,a} \\
D^2\Phi^I_a| & \equiv & F^I_a & \Gamma_{\alpha\,ab}| & \equiv & \chi_{\alpha\,ab} \\
\frac{1}{2}\,D^\alpha\Gamma_{\alpha\,ab}| & \equiv & B_{ab} & D^2\Gamma_{\alpha\,ab}| & \equiv & 2\lambda_{\alpha\,ab} - i\,\partial_\alpha{}^\beta\chi_{\beta\,ab} \\
D_\alpha\Gamma^\beta{}_{ab}| & \equiv & i\,(\gamma_\mu)_\alpha{}^\beta\,A^\mu{}_{ab} - \delta_\alpha^\beta B_{ab} & D^2\Gamma^\alpha{}_{ab}| & \equiv & 2\lambda^\alpha{}_{ab} + i\,\partial^\alpha{}_\beta\chi^\beta{}_{ab} \\
D^\alpha\Gamma_{\beta\,ab}| & \equiv & i\,(\gamma_\mu)^\alpha{}_\beta\,A^\mu{}_{ab} + \delta^\alpha_\beta B_{ab} & \frac{1}{2}D^\beta D_\alpha\Gamma_{\beta\,ab}| & \equiv & \lambda_{\alpha\,ab}
\end{array}$$

In the analog of the Wess-Zumino procedure we can gauge away B and χ component fields. Our spacetime signature is $(-,+,+)$. The purely immaginary totally antisymmetric symbol $C_{\alpha\beta}$ is used to raise and lower spinor indexes according to the $\searrow$ convention

$$\begin{array}{rcll}
C_{\alpha\beta} & = & -C_{\beta\alpha} = -C^{\alpha\beta} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} & C_{\alpha\beta}C^{\gamma\delta} = \delta_{[\alpha}^\gamma\delta_{\beta]}^\delta \\
\psi_\alpha & = & \psi^\beta C_{\beta\alpha} \qquad\qquad \psi^\alpha = C^{\alpha\beta}\psi_\beta & \psi^2 = \frac{1}{2}\psi^\alpha\psi_\alpha
\end{array}$$

We represent vectors in spinor notation as symmetric matrices:

$$\begin{array}{rcccccccl}
(\gamma^0)_{\alpha\beta} & = & -(\gamma^0)^{\alpha\beta} & = & -(\gamma_0)_{\alpha\beta} & = & (\gamma_0)^{\alpha\beta} & = & \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\
(\gamma^1)_{\alpha\beta} & = & (\gamma^1)^{\alpha\beta} & = & (\gamma_1)_{\alpha\beta} & = & (\gamma_1)^{\alpha\beta} & = & \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\
(\gamma^2)_{\alpha\beta} & = & (\gamma^2)^{\alpha\beta} & = & (\gamma_2)_{\alpha\beta} & = & (\gamma_2)^{\alpha\beta} & = & \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}
\end{array}$$

$$(\gamma^\mu)_{\alpha\beta}V_\mu = \begin{pmatrix} V_0 + V_1 & V_2 \\ V_2 & V_0 - V_1 \end{pmatrix}$$

Then the following relations hold:

$$
\begin{array}{rclrcl}
\partial^{\alpha\beta} & = & (\gamma_\mu)^{\alpha\beta}\,\partial^\mu & \partial^\mu & = & \frac{1}{2}\,(\gamma^\mu)_{\alpha\beta}\,\partial^{\alpha\beta} \\
x^{\alpha\beta} & = & \frac{1}{2}\,(\gamma_\mu)^{\alpha\beta}\,x^\mu & x^\mu & = & (\gamma^\mu)_{\alpha\beta}\,x^{\alpha\beta} \\
A^{\alpha\beta} & = & \frac{1}{\sqrt{2}}\,(\gamma_\mu)^{\alpha\beta}\,A^\mu & A^\mu & = & \frac{1}{\sqrt{2}}\,(\gamma^\mu)_{\alpha\beta}\,A^{\alpha\beta} \\
\partial_{\alpha\beta}x^{\gamma\rho} & = & \frac{1}{2}\,\delta_{(\alpha}^{\ \gamma}\delta_{\beta)}^{\ \rho} & \partial_\alpha\theta^\beta & = & \delta_\alpha^{\ \beta} \\
(\gamma_\mu)^{\alpha\beta}(\gamma_\nu)_{\alpha\beta} & = & 2\eta_{\mu\nu} & (\gamma^\mu)_{\alpha\beta}(\gamma_\mu)^{\gamma\rho} & = & \delta_\alpha^{\ \gamma}\delta_\beta^{\ \rho} + \delta_\alpha^{\ \rho}\delta_\beta^{\ \gamma} \\
(\gamma^\mu)^{\alpha\beta}(\gamma^\nu)_{\beta\rho} & = & \eta^{\mu\nu}\delta^\alpha_\rho + i\epsilon^{\mu\nu\sigma}(\gamma_\sigma)^\alpha_{\ \ \rho} & (\gamma^\mu)^\rho_{\ \rho} & = & 0 \\
(\gamma^\mu)^{\alpha\beta}(\gamma^\nu)_\beta^{\ \rho}(\gamma^\sigma)_{\rho\alpha} & = & 2i\epsilon^{\mu\nu\sigma} & \epsilon^{012} & = & 1
\end{array}
$$

We conclude with some useful relations for three dimensional D-algebra computations:

$$
\begin{array}{rclrcl}
D^\alpha D_\beta & = & i\,\partial^\alpha_\beta + \delta^\alpha_\beta\,D^2 & D_\alpha D^\beta & = & i\,\partial_\alpha^{\ \beta} - \delta_\alpha^{\ \beta}\,D^2 \\
D^2 D_\alpha & = & -i\,\partial_\alpha^{\ \beta}\,D_\beta & D_\alpha D^2 & = & i\,\partial_\alpha^{\ \beta}\,D_\beta \\
D^2 D^\alpha & = & i\,\partial^\alpha_\beta\,D^\beta & D^\alpha D^2 & = & -i\,\partial^\alpha_\beta\,D^\beta \\
D^\alpha D_\beta D_\alpha & = & 0 & \partial^{\alpha\beta}\partial_{\gamma\beta} & = & \delta^\alpha_\gamma\,\Box \\
D^2 D^2 & = & \Box & D^\alpha D^2 D_\alpha & = & -2\,\Box \\
D^2 D_\alpha D^2 & = & -\,\Box\,D_\alpha & \Box & = & \partial^\mu\partial_\mu \;=\; \frac{1}{2}\partial^{\alpha\beta}\partial_{\alpha\beta}
\end{array}
$$

B Octonionic conventions

In the octonion algebra we can choose a basis of elements

$$\Big\{1,\, e_i\Big\} \qquad\qquad i = 1,..,7$$

such that

$$e_i e_j = c_{ijk}\, e_k - \delta_{ij}$$

where the tensor c_{ijk} is totally antisymmetric with non vanishing entries:

$$c_{123} = c_{147} = c_{165} = c_{246} = c_{257} = c_{354} = c_{367} = 1$$

We can also introduce the seven dimensional dual of the structure constants:

$$c_{ijkl} = \frac{1}{6}\, \epsilon_{ijklmno}\, c^{mno}$$

Combining these two objects one can construct an $SO(7)$ invariant tensor $C_{IJKL\ \ I,J,K,L=1,...,8}$ which is self dual in 8 dimensions, by taking:

$$C_{ijk8} = c_{ijk} \qquad\qquad C_{ijkl} = c_{ijkl}$$

Octonionic structure constants can be used to construct $SO(8)$ gamma matrices. For instance a suitable representation of the triality tensor that enters BL susy transformations is given by:

$$\begin{array}{rcl} (\Gamma^i)_{A\dot{A}} & = & c^i{}_{A\dot{A}} + \delta_{8\dot{A}}\delta_{Ai} - \delta_{8A}\delta_{\dot{A}i} \qquad\qquad i = 1,...,7 \qquad A, \dot{A} = 1,...,8 \\ (\Gamma^8)_{A\dot{A}} & = & \delta_{A\dot{A}} \qquad\qquad\qquad\qquad\qquad\quad c^i_{8\dot{A}} = c^i_{A8} = 0 \end{array}$$

Defining $\hat{\Gamma}^I_{\dot{A}A} = \left(\Gamma^T\right)^I_{\dot{A}A}$ it is easy to see that $\Gamma^I\hat{\Gamma}^J + \Gamma^J\hat{\Gamma}^I = 2\delta^{IJ}$ and therefore we can write down 16x16 matrices satisfying the clifford algebra

$$\gamma^I = \left(\begin{array}{cc} 0 & \Gamma^I{}_{A\dot{A}} \\ \hat{\Gamma}^I{}_{\dot{A}A} & 0 \end{array}\right), \qquad\qquad \gamma^I\gamma^J + \gamma^J\gamma^I = 2\delta^{IJ}\,.$$

With the above definitions it can be shown that

$$\Gamma^{IJ}_{AB} = \frac{1}{2}\left(\Gamma^I_{A\dot{A}}\hat{\Gamma}^J_{\dot{A}B} - \Gamma^J_{A\dot{A}}\hat{\Gamma}^I_{\dot{A}B}\right) = C^{IJ}{}_{AB} + \delta^I_A\delta^J_B - \delta^I_B\delta^J_A\,.$$

so that the antisymmetrized product of Γ's of the scalar-fermion interaction can be written by means of the C_{IJKL} tensor.

References

[1] S. de Haro and A. C. Petkou, "Holographic Aspects of Electric-Magnetic Dualities," arXiv:0710.0965 [hep-th].

[2] J. Bagger and N. Lambert, "Modeling multiple M2's," Phys. Rev. D **75**, 045020 (2007) [arXiv:hep-th/0611108].

[3] J. Bagger and N. Lambert, "Gauge Symmetry and Supersymmetry of Multiple M2-Branes," Phys. Rev. D **77**, 065008 (2008) [arXiv:0711.0955 [hep-th]].

[4] J. Bagger and N. Lambert, "Comments On Multiple M2-branes," JHEP **0802** (2008) 105 [arXiv:0712.3738 [hep-th]].

[5] A. Gustavsson, "Algebraic structures on parallel M2-branes," arXiv:0709.1260 [hep-th].

[6] A. Gustavsson, "Selfdual strings and loop space Nahm equations," JHEP **0804** (2008) 083 [arXiv:0802.3456 [hep-th]].

[7] S. Mukhi and C. Papageorgakis, "M2 to D2," JHEP **0805**, 085 (2008) [arXiv:0803.3218 [hep-th]].

[8] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, "N = 8 Superconformal Chern–Simons Theories," JHEP **0805** (2008) 025 [arXiv:0803.3242 [hep-th]].

[9] D. S. Berman, L. C. Tadrowski and D. C. Thompson, "Aspects of Multiple Membranes," arXiv:0803.3611 [hep-th].

[10] M. van Raamsdonk, "Comments on the Bagger-Lambert theory and multiple M2-branes", arXiv:0803.3803 [hep-th].

[11] A. Morozov, "On the Problem of Multiple M2 Branes," JHEP **0805** (2008) 076, arXiv:0804.0913 [hep-th].

[12] N. Lambert and D. Tong, "Membranes on an Orbifold", arXiv:0804.1114 [hep-th].

[13] J. Distler, S. Mukhi, C. Papageorgakis and M. Van Raamsdonk, "M2-branes on M-folds," JHEP **0805**, 038 (2008) arXiv:0804.1256 [hep-th].

[14] U. Gran, B. E. W. Nilsson and C. Petersson, "On relating multiple M2 and D2-branes," arXiv:0804.1784 [hep-th].

[15] P. M. Ho, R. C. Hou and Y. Matsuo, "Lie 3-Algebra and Multiple M2-branes," arXiv:0804.2110 [hep-th].

[16] E. A. Bergshoeff, M. de Roo and O. Hohm, "Multiple M2-branes and the Embedding Tensor," arXiv:0804.2201 [hep-th].

[17] J. Gomis, A. J. Salim and F. Passerini, "Matrix Theory of Type IIB Plane Wave from Membranes," arXiv:0804.2186 [hep-th].

[18] K. Hosomichi, K. M. Lee and S. Lee, "Mass-Deformed Bagger-Lambert Theory and its BPS Objects," arXiv:0804.2519 [hep-th].

[19] G. Papadopoulos, "M2-branes, 3-Lie Algebras and Plucker relations," JHEP **0805** (2008) 054 [arXiv:0804.2662 [hep-th]].

[20] D. Gaiotto and E. Witten, "Janus Configurations, Chern-Simons Couplings, And The Theta-Angle in N=4 Super Yang-Mills Theory", arXiv:0804.2907 [hep-th].

[21] H. Shimada, "β-deformation for matrix model of M-theory," arXiv:0804.3236 [hep-th].

[22] G. Papadopoulos, "On the structure of k-Lie algebras," arXiv:0804.3567 [hep-th].

[23] P. M. Ho and Y. Matsuo, "M5 from M2," arXiv:0804.3629 [hep-th].

[24] J. Gomis, G. Milanesi and J. G. Russo, "Bagger-Lambert Theory for General Lie Algebras," arXiv:0805.1012 [hep-th].

[25] S. Benvenuti, D. Rodriguez-Gomez, E. Tonni and H. Verlinde, "N=8 superconformal gauge theories and M2 branes," arXiv:0805.1087 [hep-th].

[26] P. M. Ho, Y. Imamura and Y. Matsuo, "M2 to D2 revisited," arXiv:0805.1202 [hep-th].

[27] A. Morozov, "From Simplified BLG Action to the First-Quantized M-Theory," arXiv:0805.1703 [hep-th].

[28] Y. Song, "Mass Deformation of the Multiple M2 Branes Theory," arXiv:0805.3193 [hep-th].

[29] I. Jeon, J. Kim, N. Kim, S. W. Kim and J. H. Park, "Classification of the BPS states in Bagger-Lambert Theory," arXiv:0805.3236 [hep-th].

[30] C. Krishnan and C. Maccaferri, "Membranes on Calibrations," arXiv:0805.3125 [hep-th].

[31] K. Hosomichi, K. M. Lee, S. Lee, S. Lee and J. Park, "N=4 Superconformal Chern-Simons Theories with Hyper and Twisted Hyper Multiplets," arXiv:0805.3662 [hep-th].

[32] S. Banerjee and A. Sen, "Interpreting the M2-brane Action," arXiv:0805.3930 [hep-th].

[33] H. Lin, "Kac-Moody Extensions of 3-Algebras and M2-branes," arXiv:0805.4003 [hep-th].

[34] J. H. Park and C. Sochichiu, "Single M5 to multiple M2: taking off the square root of Nambu-Goto action," arXiv:0806.0335 [hep-th].

[35] A. Gustavsson, "One-loop corrections to Bagger-Lambert theory," arXiv:0805.4443 [hep-th].

[36] M. A. Bandres, A. E. Lipstein and J. H. Schwarz, "Ghost-Free Superconformal Action for Multiple M2-Branes," arXiv:0806.0054 [hep-th].

[37] F. Passerini, "M2-Brane Superalgebra from Bagger-Lambert Theory," arXiv:0806.0363 [hep-th].

[38] J. Figueroa-O'Farrill, "Lorentzian Lie n-algebras," arXiv:0805.4760 [math.RT].

[39] O. Aharony, O. Bergman, D. L. Jafferis and J. Maldacena, "N=6 superconformal Chern-Simons-matter theories, M2-branes and their gravity duals," arXiv:0806.1218 [hep-th].

[40] J. Figueroa-O'Farrill, P. de Medeiros and E. Mendez-Escobar, "Lorentzian Lie 3-algebras and their Bagger-Lambert moduli space," arXiv:0805.4363 [hep-th].

[41] J. Gomis, D. Rodriguez-Gomez, M. Van Raamsdonk and H. Verlinde, "The Superconformal Gauge Theory on M2-Branes," arXiv:0806.0738 [hep-th].

[42] M. Benna, I. Klebanov, T. Klose and M. Smedback, "Superconformal Chern-Simons Theories and AdS_4/CFT_3 Correspondence", arXiv:0806.1519 [hep-th].

[43] B. Ezhuthachan, S. Mukhi and C. Papageorgakis, "D2 to D2," arXiv:0806.1639 [hep-th].

[44] L. V. Avdeev, D. I. Kazakov and I. N. Kondrashuk, "Renormalizations in supersymmetric and nonsupersymmetric nonAbelian Chern-Simons field theories with matter," Nucl. Phys. B **391**, 333 (1993).

[45] L. V. Avdeev, G. V. Grigorev and D. I. Kazakov, "Renormalizations in Abelian Chern-Simons field theories with matter," Nucl. Phys. B **382**, 561 (1992).

[46] S.J. Gates, M.T. Grisaru, M. Rocek and W. Siegel, "Superspace", Benjamin-Cummings, Reading, MA, 1983. *Second printing*: Front. Phys. **58** (1983) 1 [arXiv:hep-th/0108200].

[47] B. de Wit and H. Nicolai, "The Parallelizing S(7) Torsion In Gauged N=8 Supergravity," Nucl. Phys. B **231** (1984) 506.

[48] M. Gunaydin and S. V. Ketov, "Seven-Sphere and the Exceptional N=7 and N=8 Superconformal Algebras," Nucl. Phys. B **467** (1996) 215 [arXiv:hep-th/9601072].

[49] I. Bakas, E. G. Floratos and A. Kehagias, "Octonionic gravitational instantons," Phys. Lett. B **445** (1998) 69 [arXiv:hep-th/9810042].

[50] A. Bilal, J. P. Derendinger and K. Sfetsos, "(Weak) G(2) holonomy from self-duality, flux and supersymmetry," Nucl. Phys. B **628** (2002) 112 [arXiv:hep-th/0111274].

[51] G. Ferretti, P. Salomonson and D. Tsimpis, "D-brane probes on G(2) orbifolds," JHEP **0203** (2002) 004 [arXiv:hep-th/0111050].